arXiv:hep-th/9210086v1 16 Oct 1992

GENERALIZED XYZ MODEL ASSOCIATED TO SKLYANIN ALGEBRA[*]

Takashi Takebe

Department of Mathematical Sciences
the University of Tokyo
Hongo 7-3-1, Bunkyo-ku, Tokyo 113, Japan

Abstract. The free energy of a lattice model, which is a generalization of the Heisenberg XYZ model with the higher spin representation of the Sklyanin algebra, is calculated by the generalized Bethe Ansatz of Takhtajan and Faddeev.

§0. Introduction.

The XYZ model, a quantum chain system with Hamiltonian

$$H = -\frac{1}{2}\sum_{n=1}^{N}(J_x\sigma_n^x\sigma_{n+1}^x + J_y\sigma_n^y\sigma_{n+1}^y + J_z\sigma_n^z\sigma_{n+1}^z),$$

was first solved by Baxter in early 70's by means of the Bethe Ansatz. This work was interpreted from the view point of the quantum spectral transformation method (QSTM) by Takhtajan and Faddeev in 1979 [TF]. Their starting point is the fundamental relation of the L operators

$$R^{12}(\lambda-\mu)L^{13}(\lambda)L^{23}(\mu) = L^{23}(\mu)L^{13}(\lambda)R^{12}(\lambda-\mu), \tag{0.1}$$

where

$$L(\lambda) = \sum_{a=0}^{3} W_a^L(\lambda)\sigma^a\otimes\sigma^a, \tag{0.2}$$

$$W_0^L(\lambda) = \frac{\theta_{11}(\lambda)}{\theta_{11}(\eta)}, W_1^L(\lambda) = \frac{\theta_{10}(\lambda)}{\theta_{10}(\eta)}, W_2^L(\lambda) = \frac{\theta_{00}(\lambda)}{\theta_{00}(\eta)}, W_3^L(\lambda) = \frac{\theta_{01}(\lambda)}{\theta_{01}(\eta)},$$

σ^a's are Pauli matrices, $\theta_{ab}(\lambda) = \theta_{ab}(\lambda;\tau)$ (we follow the notations of [M] for the theta functions), and the R matrix is the Baxter's R matrix defined by

$$R(\lambda) = \sum_{a=0}^{3} W_a^R(\lambda)\sigma^a\otimes\sigma^a, W_a^R(\lambda) := W_a^L(\lambda+\eta).$$

---

[*] talk given at the XXI Differential Geometry Methods in Theoretical Physics, Tianjin, China 5-9 June 1992 ; hep-th/9210086

Typeset by $\mathcal{A}_{\mathcal{M}}\mathcal{S}$-TeX

Sklyanin found the algebraic structure underlying the above relation, replacing the L operator by

$$L(\lambda) = \sum_{a=0}^{3} W_a^L(\lambda)\sigma^a \otimes S^a,$$

and writing down the relations among S^a's equivalent to (0.1) [Sk1]. The algebra Q generated by S^a's is called the Sklyanin algebra. He also constructed series of representations in [Sk2], among which we use the spin l representation ρ_l : $Q \to \mathrm{End}(V_l)$, $\dim V_l = 2l+1$. The algebra Q is a deformation of the universal enveloping algebra $U(sl(2,\mathbb{C}))$, and ρ_l is corresponding deformation of the spin l representation of $sl(2,\mathbb{C})$.

As the solvability of the XYZ (or 8 vertex) model comes from the fundamental relation (0.1), it is natural to consider the generalization of this model by using the above representations of the Sklyanin algebra, and calculate its free energy.

§**1. Description of the model.** Now we construct a generalization of the XYZ model by replacing the L operators of the form (0.2) with

$$(1.1)\qquad \begin{aligned} L_n(\lambda) &= \sum_{a=0}^{3} W_a^L(\lambda)\,\sigma^a \otimes \rho_l^{(n)}(S^a), \\ \rho_l^{(n)}(S^a) &= 1 \otimes \cdots \otimes 1 \otimes \underset{n-\text{th component}}{\rho_l(S^a)} \otimes 1 \otimes \cdots 1, \end{aligned}$$

which act on the space $\mathbb{C}^2 \otimes V_l^{\otimes N}$. The transition matrix of the higher spin generalization of the 8 vertex model is defined by:

$$T(\lambda) = L_N(\lambda) \cdots L_2(\lambda) L_1(\lambda).$$

The transition matrix of the corresponding chain model is

$$t(\lambda) = \mathrm{tr}_{\mathbb{C}^2}\, T(\lambda),$$

which acts on the space $V_l^{\otimes N}$. The free energy of the model under consideration is, by definition, the Perron-Frobenius eigenvalue of the transfer matrix $t(\lambda)$, Λ_N, or its thermodynamic limit,

$$f = \lim_{N\to\infty} \frac{1}{N} \ln \Lambda_N.$$

Similarly to the choice of [TF], we assume that the elliptic modulus τ is of the form $\tau = i/t$, where $t > 0$ and the parameters η, λ belong to the domain $0 < \lambda + \eta < \max(\eta, 2l\eta) < \frac{1}{2}$ (antiferromagnetic region).

In the previous paper [Take], the author showed that the generalized Bethe Ansatz of Takhtajan and Faddeev is applicable to this model. According to this result, if $M = Nl$ parameters, $\lambda_1, \ldots \lambda_M$, satisfy the Bethe equation,

$$(1.2)\qquad \left(\frac{\theta_{11}(\lambda_j - 2l\eta)}{\theta_{11}(\lambda_j + 2l\eta)}\right)^N = \prod_{k=1, k\neq j}^{M} \frac{\theta_{11}(\lambda_k - \lambda_j - 2\eta)}{\theta_{11}(\lambda_k - \lambda_j + 2\eta)},$$

for $j = 1, \ldots, M$, then the transfer matrix $t(\lambda)$ has an eigenvalue
(1.3)

$$(2\theta_{11}(\lambda - 2l\eta))^N \prod_{k=1}^{M} \frac{\theta_{11}(\lambda - \lambda_k + 2\eta)}{\theta_{11}(\lambda - \lambda_k)} + (2\theta_{11}(\lambda + 2l\eta))^N \prod_{k=1}^{M} \frac{\theta_{11}(\lambda + \lambda_k + 2\eta)}{\theta_{11}(\lambda - \lambda_k)}.$$

§2. Thermodynamic limit. Now we take the thermodynamic limit $N \to \infty$. From the results of the XXX and XXY models [Takh], [B] and [S], we assume that the following *string hypothesis* holds also for this model: assymptotically as $N \to \infty$, the solutions of the Bethe equation (1.2) corresponding to the ground state cluster in complexes, so called $2l$-strings:

$$\{\lambda_k^0 + 2\alpha\eta |\, \alpha = -l + \tfrac{1}{2}, \dots , l + \tfrac{1}{2}\}_{k=1,\dots,N/2}$$

(assuming that N is even), and centres of these strings λ_k^0 distribute in a purely imaginary interval $\{ix | -1/2t < x < 1/2t\}$ and have a smooth density $\rho(\lambda')d\lambda'$ as $N \to \infty$ ($\lambda' = \lambda/\tau$).

Under these hypotheses, the Bethe Ansatz equation (1.2) reduces to an integral equation

$$\sum_{\alpha=0}^{2l-1} \Psi(\lambda, (2\alpha+1)\eta) = \tag{2.1}$$

$$= -2\pi\rho(\lambda') + \int_{-1/2}^{1/2} \Big(\sum_{\beta=1}^{2l-1} \Psi(\lambda - \mu, 2\beta\eta) + \sum_{\beta=0}^{2l-1} \Psi(\lambda - \mu, 2(\beta+1)\eta)\Big)\rho(\mu')\, d\mu',$$

where $\lambda' = \lambda/\tau$, $\mu' = \mu/\tau$ and

$$\Psi(\lambda, \eta) = \frac{\tau}{i}\frac{d}{d\lambda} \ln \frac{\theta_{11}((\lambda-\eta)/\tau; -1/\tau)}{\theta_{11}((\lambda+\eta)/\tau; -1/\tau)}.$$

We can solve this equation, expanding $\rho(\lambda')$ into Fourier series, and obtain

$$\rho(\lambda') = \sum_{n\in\mathbb{Z}} \frac{e^{2\pi i n\lambda'}}{2\cos 2\pi n\eta'}. \tag{2.2}$$

Using this density of the solutions of the Bethe equation, we can take the limit of the eigenvalue (1.3) and obtain the free energy of the model $f = \lim_{N\to\infty} \ln \Lambda_N/N$ as

$$e^f \propto \theta_{11}(\lambda + 2l\eta) e^{2\pi t(\lambda-\eta)(2l\eta - \frac{1}{2})} \times \tag{2.3}$$

$$\times \exp \sum_{n=1}^{\infty} \frac{\sinh 2\pi n t(2l\eta - \frac{1}{2})\, \sinh 2\pi n t(\lambda - \eta)}{n \sinh \pi n t\, \cosh 2\pi n t\eta} =$$

$$= \theta_{11}(\lambda + 2l\eta)\, q^{-(\lambda-\eta)(2l\eta - \frac{1}{2})} \frac{\sigma(\lambda; \eta, l|q)}{\sigma(-\lambda; \eta, l|q)},$$

where $q = \exp(-2\pi t)$ and

$$\sigma(\lambda; \eta, l|q) = \prod_{k=0}^{\infty} \frac{\Gamma_q(\lambda + 4\eta k + 2(l+1)\eta)\, \Gamma_q(\lambda + 4\eta k - 2l\eta + 1)}{\Gamma_q(\lambda + 4\eta k + 2l\eta)\, \Gamma_q(\lambda + 4\eta k - 2(l-1)\eta + 1)},$$

$$\Gamma_q(x) = \frac{(q;q)_\infty}{(q^x;q)_\infty}(1-q)^{1-x}, \qquad (a;q)_\infty = \prod_{n=0}^{\infty}(1 - aq^n).$$

This result is consistent with that of [TF] which corresponds to the case $l = \frac{1}{2}$.

Details and the further results concerning the thermodynamic properties and the excited states of this model will be published in the forthcoming paper.

The author expresses his great thanks to Professors E. K. Sklyanin, H. J. de Vega, M. Wadati, E. Date, Kimio Ueno, M. Noumi and Dr. T. Deguchi for their comments, advices, interests and encouragement.

REFERENCES

[B] Babujian H.M., Phys.lett.A **90** (1982), 479–482.

[M] Mumford, D., *Tata Lectures on Theta I*, Birkhäuser, 1982.

[S] Sogo K., Phys.lett.A **104** (1984), 51–54.

[Sk1] Sklyanin E.K., Func. Anal. Appl. **16-4** (1982), 263–270.

[Sk2] ______, Func. Anal. Appl. **17-4** (1983), 273–284.

[Take] Takebe T., J.Phys.A **25** (1992), 1071–1083.

[Takh] Takhtajan L.A., Phys.lett.A **87** (1982), 479–482.

[TF] Takhtajan L.A., Faddeev L.D., Uspekhi Mat. Nauk **34:5** (1979), 13–63; English transl. in Russ. Math. Surv. **34** (1979), 11–68.